ELSEVIER

Toxicology 117 (1997) 247–248

TOXICOLOGY

Volume contents

Volume 117 (1997)

Structural basis for the haemotoxicity of dapsone: the importance of the sulphonyl group
In a series of analogues dapsone itself is the most potent methemoglobinogenic agent
R. Mahmud, M.D. Tingle, J.L. Maggs, M.T.D. Cronin, J.C. Dearden, B.K. Park (UK) 1–11

Interaction of methadone with substrates of human hepatic cytochrome P450 3A4
Inhibition by methadone is fairly specific to P450 3A4. Fluvoxemine inhibits methadone N-demethylation
C. Iribarne, Y. Dréano, L.G. Bardou, J.F. Ménez, F. Berthou (France) 13–23

A pharmacokinetic model describing pulsatile uptake of orally-administered carbon tetrachloride
Estimates of absorption rates and bioavailabilities are provided by the model
G. Semino, P. Lilly, M.E. Andersen (Italy, USA) 25–33

Membrane toxicity of opioids measured by protozoan motility
In high concentrations opioids may have a receptor-independent membrane action
C. Wu, C.H. Fry, J.A. Henry (UK) 35–44

Biphasic effects of chromium compounds on catecholamine secretion from bovine adrenal medullary cells
Chromium compounds have diverse effects on catecholamine secretion, possibly by interference with intracellular calcium
P.-S. Liu, M.-K. Lin (Peoples Republic of China) 45–53

Oxidative liver DNA damage in rats treated with pesticide mixtures
Of 15 tested pesticides only diphenylamine and chlorothalonil increase the ratio of 8-hydroxy-deoxyguanosine versus 2-deoxyguanosine
M. Lodovici, C. Casalini, C. Briani, P. Dolara (Italy) 55–60

Effects of triethyl lead administration on the expression of glutathione S-transferase isoenzymes and quinone reductase in rat kidney and liver
Glutathione-s-transferase expression and quinone oxidoreductase activity do not appear to be co-regulated
D.A. Daggett, E.F. Nuwaysir, S.A. Nelson, L.S. Wright, S.E. Kornguth, F.L. Siegel (USA). 61–71

The effect of aromatic fluorine substitution on the nephrotoxicity and metabolism of *N*-(3,5-dichlorophenyl)succinimide in Fischer 344 rats
DFPS and its oxidative metabolites are mild nephrotoxicants that produce reversible changes
G.L. Kellner-Weibel, A. K. Nyarko, R. Tchao, C.M. Henesey, P. J. Harvison (USA, Ghana) 73–83

Book review 85
Instructions to Authors 87–88

Copper ions differ from other thiol reactive metal ions in their effects on the concentration and redox status of thiols in HeLa cell cultures
Cu^{2+} is the most efficient ion to disturb thiol homeostasis
B. Hultberg, A. Andersson, A. Isaksson (Sweden) 89–97

Lack of correlation between the inducibility of metallothionein mRNA and metallothionein protein in cadmium-exposed rodents
Cadmium toxicity and carcinogenicity depends upon Cd distribution, inducibility of the metallothionein gene and organ levels of metallothionein protein
R.R. Misra, K.A. Crance, R.M. Bare, M.P. Waalkes (USA) 99–109

G-Protein(s), $G_{\alpha q}/G_{\alpha 11}$, in the olfactory neuroepithelium of the channel catfish (*Ictalurus punctatus*) is altered by the herbicide, dichlobenil
The herbicide may be a powerful olfactory toxicant in fish
I. Andreini, C. DellaCorte, L.C. Johnson, S. Hughes, D.L. Kalinoski (USA, Italy) 111–122
N-Deacetyl ketoconazole-induced hepatotoxicity in a primary culture system of rat hepatocytes
Flavin-containing monoxygenases may mediate toxicity
R.J. Rodriguez, D. Acosta, Jr (USA). 123–131
Tyrosine prevents capsaicin-induced protein synthesis inhibition in cultured cells
Tyrosine fairly specifically reduces cytotoxicity of capsaicin
C. Cochereau, D. Sanchez, E.E. Creppy (France). 133–139
Characterization of Fe^{2+} and Fe^{3+} transport by iron-loaded cardiac myocytes
In iron overload, both redox states of iron contribute to cardiac iron accumulation
J.G. Parkes, N.F. Olivieri, D.M. Templeton (Canada) 141–151
Lead perturbs the regulation of spontaneous release of tissue plasminogen activator and plasminogen activator inhibitor-1 from vascular smooth muscle cells and fibroblasts in culture
An involvement of mainly intracellular calcium-independent mechanisms is suggested
C. Yamamoto, A. Miyamoto, M. Sakamoto, T. Kaji, H. Kozuka (Japan) 153–161
Cardiovascular collapse through non-cholinergic mechanism after intravenous injection of *N*-methylcarbamate insecticide in rabbits
Differences in cardiovascular response may depend on a balance between cholinergic and non-cholinergic activity of an inhibitor
H. Futagawa, Y. Kakinuma, H. Takahashi (Japan) 163–170
Protection against cocaine and combined cocaine–ethanol toxicities in mice by imidazobenzodiazepine Ro 15-4513
The effectiveness of the drug is documented
T. Hayase, Y. Yamamoto, K. Yamamoto (Japan) 171–182
Association of HSP73 with the acquired resistance to uranyl acetate-induced acute renal failure
A possible protective mechanism
S. Mizuno, K. Fujita, R. Furuy, A. Hishid, H. Ito, Y. Tashim, H. Kumagai (Japan). 183–191
Inhibitory effect of lead on the repair of wounded monolayers of cultured vascular endothelial cells
No effect was found with other heavy metals such as Hg, Bi, Co, Mn or Ni
Y. Fujiwara, T. Kaji, S. Sakurai, Michiko Sakamoto, H. Kozuka (Japan) 193–198
Regression of morphological alterations and oxidative stress-related parameters after acute lindane-induced hepatotoxicity in rats
A time-course study
V.B.C. Junqueira, O.R. Koch, A.C.M. Arisi, A.P. Fuzaro, L.A. Azzalis, S.B.M. Barros, A. Cravero, S. Farré, L.A. Videla (Brasil, Argentina, Chile). 199–205
Buthionine sulfoximine (BSO) and *N*-(3,5-dichlorophenyl)succinimide nephrotoxicity: temporal aspects of BSO administration and BSO effects on renal transport systems
BSO attenuates nephrotoxicity through glutathione depletion
G.O. Rankin, K.W. Beers, V.J. Teets, D.W. Nicoll, D.K. Anestis (USA). 207–217

Immunotoxicology
Effect of saponin on the transmucosal passage of β-lactoglobulin across the proximal small intestine of normal and β-lactoglobulin-sensitised rats.
Exposure to saponin increased the transmucosal uptake of β-lactoglobulin
J.M. Gee, J.M. Wal, K. Miller, H. Atkinson, F. Grigoriadou, M.V.W. Wijnands, A.H. Penninks, G. Wortley, I.T. Johnson (UK, France, The Netherlands). 219–228
Local lymph node activation and IgE responses in Brown Norway and Wistar rats after dermal application of sensitizing and non-sensitizing chemicals
Trimetallic anhydride, but neither dinitrochlorobenzene, formaldehyde or methylsalicylate, produced an increase in serum IgE levels following topical exposure at Brown Norway rats
J.H.E. Arts, S.C.M. Dröge, S. Spanhaak, N. Bloksma, A.H. Penninks, C.F. Kuper (The Netherlands) 229–237

Announcement 239–240
Author Index. 241–242
Subject Index. 243–246
Volume Contents 247–249